

Tarbiat Modares University Press  
Entomological Society of Iran

Short paper

Faunistic

<https://doi.org/10.52547/jibs.9.2.303>

ISSN: 2423-8112

<https://zoobank.org/urn:lsid:zoobank.org:F63D09F0-1D81-4F99-97E6-C4F33A99C3B0>

## *Diglyphus anadolucus* Doğanlar, 1982 (Hym., Eulophidae) a primary larval ectoparasitoid of Anthomyiidae (Diptera), with a checklist of *Diglyphus* species in Iran

**Younes Karimpour**

Department of Plant Protection, Faculty of Agriculture, Urmia University, Urmia – Iran.

✉ [y.karimpour@urmia.ac.ir](mailto:y.karimpour@urmia.ac.ir) <https://orcid.org/0000-0003-2468-8367>**Majid Jafarlu**

Department of Plant Protection, Faculty of Agriculture, Urmia University, Urmia – Iran.

✉ [m.jafarlu@urmia.ac.ir](mailto:m.jafarlu@urmia.ac.ir) <https://orcid.org/0000-0002-5829-8781>**Hossein Lotfalizadeh**

Insects Taxonomy Research Department, Iranian Research Institute of Plant Protection (IRIPP), Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran.

✉ [h.lotfalizadeh@areeo.ac.ir](mailto:h.lotfalizadeh@areeo.ac.ir) <https://orcid.org/0000-0002-7927-819X>

**ABSTRACT.** This is the first case of host record for parasitoid wasps of the genus *Diglyphus* Walker (Hymenoptera, Eulophidae) from the Anthomyiidae (Diptera) family. This result is based on specimens of *D. anadolucus* Doğanlar, 1982 which reared from parasitized larval specimen of *Pegomya terebrans* (Rondani) as a leaf miner of cotton thistle, *Onopordum acanthium* L. (Asteraceae). *Diglyphus anadolucus* is reported from Iran for the first time. Some biological notes and morphological characteristics of *D. anadolucus* along with related photographs are given. An updated checklist of the twelve known *Diglyphus* species in Iran together with their host associations is also provided.

**Key words:** biocontrol, leafminers, new host association, new record

**Received:**

24 December, 2022

**Accepted:**

29 January, 2022

**Published:**

05 February, 2023

**Subject Editor:**

Ehsan Rakhshani

**Citation:** Karimpour, Y., Jafarlu, M. & Lotfalizadeh, H. (2023) *Diglyphus anadolucus* Doğanlar, 1982 (Hym., Eulophidae) a primary larval ectoparasitoid of Anthomyiidae (Diptera), with a checklist of *Diglyphus* species in Iran. *Journal of Insect Biodiversity and Systematics*, 9 (2), 303–310.

## INTRODUCTION

Species of the genus *Diglyphus* Walker, 1844 (Eulophidae, Eulophinae) with worldwide distribution includes 41 known species, of which 31 species are distributed in the Palaearctic region (Noyes, 2019), and 11 species in Iran (Jafarlu et al., 2022). They are primary, solitary or gregarious ectoparasitoids on, mainly leaf mining Agromyzidae (Bouček & Askew, 1968), although, some species have also been reported on different hosts than Agromyzidae, especially on Lepidoptera: Gracillariidae, Lyonetiidae and Nepticulidae (Noyes, 2019). More recently, Jafarlu et al. (2022) reviewed the *Diglyphus* species in lucerne fields of Iran. Among them, there is no report on parasitizing leaf miners of the family Anthomyiidae.

## MATERIAL AND METHODS

The leaf samples of *Onopordum acanthium* L. (Asteraceae) infested with a leaf miner were collected along with infected leaves from the campus of Urmia University on June 5, 2020. Samples were sorted


**Corresponding author:** Karimpour, Y., E-mail: [y.karimpour@urmia.ac.ir](mailto:y.karimpour@urmia.ac.ir)

**Copyright** © 2023, Karimpour et al. This is an open access article distributed under the terms of the Creative Commons NonCommercial Attribution License (CC BY NC 4.0), which permits Share - copy and redistribute the material in any medium or format, and Adapt - remix, transform, and build upon the material, under the Attribution-NonCommercial terms.

and transferred to the laboratory, where they were reared inside the muslin-ventilated boxes for one week. The infested leaves were inspected under a stereomicroscope, for possible parasitoids. Once the parasitized larvae were detected, (Fig. 1), the leaves were carefully moved into a separate petri dish and kept under lab conditions, until the emergence of the adult parasitoids. The parasitoid specimens were transferred into 75% ethanol. For mounting, specimens were treated using the method suggested by Noyes (1982) so that dehydration was managed using alcoholic series of 75%, 85%, and 96%. Subsequently, for the drying phase to prevent specimens from shriveling, the CPD (critical point drying) method was utilized. The acetic acid bath was used because of the convenient utilization of the specimens during card-mounting. Rectangular cards with dimensions of 0.5 × 1.4 cm were used for card-mounting and water-soluble glue to fix specimens on the rectangular cards (Noyes, 1982). Morphological characters of specimens were examined under an Olympus SZH stereomicroscope. Images were taken with a Keyence digital microscope (VHX-5000) and edited in in Adobe Photoshop® CS6 software. The terminology followed Gibson (1997) and Yoder et al. (2010). Identification at generic and species levels was made using identification keys in Hansson and Navone (2017) and Doğanlar (1982). Identified species were deposited in Hayk Mirzayans Insect Museum, Tehran, Iran (HMIM).

## RESULTS

Adult specimens of ectoparasitoid wasp were emerged after 9–10 days through rearing of their larvae on leaf miner larva. The leaf miner was identified as *Pegomya terebrans* (Rondani) (Diptera, Anthomyiidae). Cotton thistle leaf miner host plants are restricted to three species of plants belonging to tribe Cynareae (Asteraceae), including Italian thistle *Carduus pycnocephalus* L. and two species of cotton thistle namely *Onopordum acanthium* L. and *O. leptolepis* DC. in Urmia environs (see Karimpour et al., 2021).



**Figure 1.** The larvae of *Diglyphus anadolucius* Doğanlar, feeding on larva of *Pegomya terebrans* (Rondani) as an ectoparasitoid.

*Taxonomic hierarchy*

Class Insecta Linnaeus, 1758

Order Hymenoptera Linnaeus, 1758

Superfamily Chalcidoidea Latreille, 1817

Family Eulophidae Westwood, 1829

Subfamily Eulophinae Westwood, 1829

Genus *Diglyphus* Walker, 1844*Diglyphus* Walker, 1844:409. Type species: *Cirrospilus chabrias* Walker, 1838:451, designated by monotypy.

**Diagnosis.** Both female and male with two funiculars and three clavomeres, males with unbranched flagellomeres. Submarginal vein smoothly joining to parastigma and having more than 3 dorsal setae. Mesoscutellum with submedian grooves, Propodeum without plicae. Gaster sessile, body medium sized, color dark and metallic.

*Diglyphus anadolucus* Doğanlar, 1982*Diglyphus anadolucus* Doğanlar, 1982:75–78. ♂♀, Holotype ♀, AUET, Türkiye. Not examined.

(Figs 2A–E)

**Material examined.** 6 ♀♀; 14–15, June 2020; ex. *Pegomya terebrans* (Rondani) (Diptera, Anthomyiidae) on *Onopordum acanthium* DC. (Asteraceae); Urmia, Urmia University campus, 37°39'14"N, 44°58'35"E, 1360 m a.s.l., leg. Y. Karimpour.

**Diagnosis – Female.** Body metallic green with golden reflections on mesoscutellum (Figs 2A, 2C, 2D) Pedicel and flagellum blackish. Scape white with anterior 0.36 in dark brown color (Fig. 2B). Femora and tibiae white, except for the hind femur which is dark brown at 1/3 its base (Fig. 2A). Antenna with first funicular 2.13× and second funicular 1.50× as long as wide; pedicel 0.63× as long as length of first flagellomere; and first funicular 1.17× as long as second funicular (Fig. 2B). Costal cell with a complete row of setae on ventral surface and with 11 setae on the dorsal surface along the terminal part of its anterior margin (Fig. 2E)

In the key provided by Hansson and Navone (2017), based on specimens obtained from Italy and Russia the first funicular of female antennae is 1.6× and second one is 1.3× as long as wide. While in the specimens of current study, they are 2.13× and 1.50× as long as wide, respectively (Fig 2B). This corresponds to the characteristics of the corresponding holotype (Doğanlar, 1982). Therefore, it is suggested that in the identification key published by Hansson and Navone (2017), the scales mentioned for first and second funiculars of female antennae be modified as follows: the first funicular 1.6–2.2× and second funicular 1.3–1.6× as long as wide.

**Distribution in Iran.** West Azarbaijan province, Urmia (present study).

**General distribution.** Iran (New record), Italy, Russia (Hansson & Navone, 2017), and Türkiye (Doğanlar, 1982).


## DISCUSSION

The genus *Diglyphus* includes 41 species worldwide (Noyes, 2019), which are mainly the ectoparasitoids of leafmining agromyzids (Diptera, Agromyzidae) (Bouček & Askew, 1968). According to the Hansson and Navone (2017) all European and Asian species of *Diglyphis* are associated with the various genera of Agromyzidae, as even the citrus leafminer, *Phyllocnistis citrella* Stainton, 1856 (Lepidoptera, Gracilariidae) (González Tirado et al., 1996). Therefore, this is the first confirmed record indicating the association of a *Diglyphus* species with the family Anthomyiidae. *Diglyphus anadolucus* was originally described from the material reared from an unknown leaf miner on Alfalfa



(Doğanlar, 1982), later it was reared from *Agromyza abiens* Zetterstedt, 1848 as a leaf miner of *Echium vulgare* L. (Boraginaceae) in Italy (Hansson & Navone, 2017). Among the 41 species of *Diglyphus* known worldwide, none of them have been reported as the primary parasitoid of Anthomyiidae (Noyes, 2019).

The plant association and primary hosts of the *Diglyphus* species reported in Iran are presented in Table 1. Including a new record for the Iranian fauna in this study, so far 12 species of the genus *Diglyphus* have been reported from Iran.



**Figure 2.** *Diglyphus anadolucus*, female: **A.** General habitus, lateral view; **B.** Head and antenna, lateral view; **C.** Mesoscutellum and propodeum, dorsal view; **D.** Head and mesosoma, dorsal view; **E.** Fore wing.

**Table 1.** List of *Diglyphus* species and their host association in Iran.

<i>Diglyphus</i> species	Collecting data			References
	Locality (province)	Host plants	Primary hosts	
<i>Diglyphus anadolucus</i> Doğanlar, 1982	West Azarbaijan	<i>Onopordum acanthium</i>	<i>Pegomya terebrans</i>	Present study
<i>D. propodealis</i> Szelenyi, 1978	West Azarbaijan	-	-	Jafarlu et al. (2022)
<i>D. chabrias</i> (Walker, 1838)	Fars	<i>Lactuca serriola</i>	<i>Liriomyza trifolii</i>	Fallahzadeh et al. (2006)
	Lorestan	-	<i>Syringopais temperatella</i>	Yefremova et al. (2007)
<i>D. crassinervis</i> Erdős, 1958	Isfahan	-	-	Ghahari & Yefremova (2013)
	Tehran	<i>Cucumis sativus</i> <i>Lycopersicon esculentum</i> <i>Citrullus lanatus</i> <i>Phaseolus vulgaris</i>	<i>Liriomyza sativae</i> <i>Liriomyza trifolii</i>	Asadi et al. (2006)
<i>D. isaea</i> (Walker, 1838)	Chaharmahal & Bakhtiari	-	-	Yefremova et al. (2007)
	East Azarbaijan	<i>Medicago sativa</i>	<i>Chromatomyia horticola</i> <i>Liriomyza sativae</i> <i>Liriomyza trifolii</i>	Lotfalizadeh et al. (2015)
	Fars	-	<i>Liriomyza trifolii</i>	Dousti et al. (2008)
	Golestan, Isfahan, Mazandaran	-	-	Ghahari & Yefremova (2013)
	Sistan & Baluchestan	<i>Triticum aestivum</i> <i>Helianthus annuus</i> <i>Silybum</i> sp. <i>Brassica rapa</i>	<i>Chromatomyia nigra</i> <i>Chromatomyia horticola</i>	Shahreki et al. (2012)
	Tehran	<i>Cucumis sativus</i> <i>Lycopersicon esculentum</i> <i>Citrullus lanatus</i> <i>Phaseolus vulgaris</i>	<i>Liriomyza sativae</i> <i>Liriomyza trifolii</i>	Asadi et al. (2006)
<i>D. minoeus</i> (Walker, 1838)	Isfahan	-	-	Yefremova et al. (2007)
<i>D. pachyneurus</i> Graham, 1963	East Azarbaijan	<i>Medicago sativa</i>	<i>Chromatomyia horticola</i> <i>Liriomyza trifolii</i>	Lotfalizadeh et al. (2015)
<i>D. poppoea</i> Walker, 1848	Fars	<i>Triticum aestivum</i>	<i>Agromyza ambigua</i>	Dousti et al. (2008)
	Golestan, Isfahan	-	-	Ghahari & Yefremova (2013)
	Sistan & Baluchestan	<i>Brassica rapa</i> <i>Cucumis sativus</i>	<i>Chromatomyia horticola</i>	Shahreki et al. (2012)
<i>D. pulchripes</i> (Crawford, 1912)	Fars	<i>Medicago sativa</i>	<i>Liriomyza trifolii</i>	Dousti et al. (2008)
<i>D. pusztensis</i> (Erdős & Novicky, 1951)	Fars	Sweeping in wheat fields	-	Hesami et al. (2010)
<i>D. sabulosus</i> Erdős, 1951	Tehran	-	-	Farahbakhsh (1961)
<i>D. scapus</i> Yefremova, 2008	Mazandaran	Sweeping in pastures	-	Hesami et al. (2006)

(-) Collected by sweeping net, and host plant/primary host is unknown.

## AUTHOR'S CONTRIBUTION

The authors confirm their contribution in the paper as follows: Y.Karimpour: Collection of the specimens, recording the biological data in Urmia. All authors confirmed their contribution in identification the specimens, sorting and deposition in the collection, preparing the diagnostic characters, preparation of the draft, corrections on the final contents of the manuscript.

## FUNDING

This research was financially supported by the Urmia University, Urmia, Iran.

## AVAILABILITY OF DATA AND MATERIAL

The specimens listed in this study are deposited in Hayk Mirzayans Insect Museum, Tehran, Iran and are available from the curator, upon request.



## ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

## CONSENT FOR PUBLICATION

Not applicable.

## CONFLICT OF INTERESTS

The authors declare that there is no conflict of interest regarding the publication of this paper.

## ACKNOWLEDGMENTS

The authors are grateful to Dr. J-Y. Rasplus (CBGP, Montpellier, France) for hosting HL and providing facilities for photography of the specimens.

## REFERENCES

- Asadi, R., Talebi, A.A., Fathipour, Y., Moharamipour, S. & Rakhshani, E. (2006) Identification of parasitoids and seasonal parasitism of the Agromyzid leafminers genus *Liriomyza* (Dip.: Agromyzidae) in Varamin, Iran. *Journal of Agricultural Science and Technology*, 8, 293–303.
- Bouček, Z. & Askew, R.R. (1968) Index of Palaearctic Eulophidae (excl. Tetrastichinae). In: Delucchi, V. & Remaudière, G. (eds) *Index of Entomophagous Insects*, 3:32, Le François, Paris, pp. 9–254.
- Doğanlar, M. (1982) A new species of *Diglyphus* Walker (Hymenoptera: Eulophidae) from eastern Anatolia. *Türkiye Bitki Koruma Dergisi*, 6, 75–79.
- Dousti, A.F., Kamali, K., Nouri-Ganbalani, G. & Ostovan, H. (2008) Report of four Hymenopteran species of Eulophidae, parasitoids of *Liriomyza trifolii* (Dip.: Agromyzidae) in Shiraz, Iran. *Journal of Entomological Society of Iran*, 27 (2), 9–10.
- Fallahzadeh, M., Naghibinejad, M., Shojaei, M. & Ostovan, H. (2006) The first report of two species of Eulophidae (Hymenoptera) from Iran. *Applied Entomology and Phytopathology*, 73 (2), 44.
- Farahbakhsh, G. (1961) *A checklist of economically important insects and other enemies of plants and agricultural products of Iran*. Ministry of Agriculture, Department of Plant Protection, Tehran. 153 p.
- Ghahari, H. & Yefremova, Z.A. (2013) A study on the Eulophidae (Hymenoptera: Chalcidoidea) from Iran. *Zoosystematica Rossica*, 22 (2), 303–310. <https://doi.org/10.31610/zsr/2013.22.2.303>
- Gibson, G.A.P. (1997) Morphology and Terminology. In: Gibson, G.A.P., Huber, J.T. & Woolley, J.B. (eds) *Annotated Keys to the Genera of Nearctic Chalcidoidea (Hymenoptera)*. National Research Council Research Press, Ottawa, Ontario, Canada. pp. 16–44.
- Gonzalez Tirado, L., Bernabe Ruiz, P. & Castano Lopez, M. (1996) Enemigos naturales autoctonos de *Phyllocnistis citrella* Stainton. (Lepidoptera. Gracillariidae, Phyllocnistinae) en la provincia de Huelva. Distribucion geografica, evolucion estacional y tasas de parasitismo. *Boletín de Sanidad Vegetal Plagas*, 22, 741–760.
- Hesami, S., Yefremova, Z.A., Ebrahimi, E. & Ostovan, H. (2006) Little known and new species of Eulophidae (Hymenoptera, Chalcidoidea) from Iran. *Entomofauna*, 27 (28), 393–402.
- Hesami, S., Ebrahimi, E., Ostovan, H. & Yefremova, Z.A. (2010) Contribution to the study of Eulophidae (Hymenoptera: Chalcidoidea) of Fars province of Iran: II - Subfamilies Entiinae and Eulophinae, with a preliminary checklist of Eulophidae species in Iran. *Plant Protection Journal*, 2, 239–253.
- Hansson, C. & Navone, P. (2017) Review of the European species of *Diglyphus* Walker (Hymenoptera: Eulophidae) including the description of a new species. *Zootaxa*, 4269 (2), 197–229. <https://doi.org/10.11646/zootaxa.4269.2.2>
- Jafarlu, M., Lotfalizadeh, H. & Karimpour, Y. (2022) Fauna of the genus *Diglyphus* (Hymenoptera: Eulophidae) in the alfalfa fields of Iran. *Journal of Entomological Society of Iran*, 43 (22), 211–219.
- Karimpour, Y., Veisi Dizaji, L., Akbarian, J. & Shahand, S. (2021) Biology of cotton thistle leaf miner, *Pegomya terebrans* (Diptera: Anthomyiidae) in Urmia region, Iran. *Journal of Entomological Society of Iran*, 41 (1), 29–40. <https://doi.org/10.22117/jesi.2021.353479.1408>

- Lotfalizadeh, H., Pourhaji, A. & Zargaran, M.R. (2015) Hymenopterous parasitoids (Hymenoptera: Braconidae, Eulophidae, Pteromalidae) of the alfalfa leafminers in Iran and their diversity. *Far Eastern Entomologist*, 288, 1–24.
- Noyes, J.S. (1982) Collecting and preserving chalcid wasps (Hymenoptera: Chalcidoidea). *Journal of Natural History*, 16, 315–334. <https://doi.org/10.1080/00222938200770261>
- Noyes, J.S. (2019) Universal Chalcidoidea database. Online Available from: <http://www.nhm.ac.uk/entomology/chalcidoids/index.html> [Accessed March 2019]
- Shahreki, Z., Rakhshani, E., Dousti, A., Ravan, S., Khajeh, N. and Taheri, S. (2012) A survey on parasitoids (Hym., Chalcidoidea, Eulophidae) of the Agromyzid leaf miners in Sistan region, Iran. In: Sarafrazi, A. (ed.) *Proceedings of 20<sup>th</sup> Iranian Plant Protection Congress*, 26–29 August 2012, Shiraz University, Shiraz, Vol I, p 103.
- Walker, F. (1838) Descriptions of British Chalcidites. *Annals and Magazine of Natural History*, 1 (6), 449–454. <https://doi.org/10.1080/00222933809496629>
- Walker, F. (1844) On the species of Chalcidites inhabiting the Arctic region. *Annals and Magazine of Natural History*, 14, 407–410. <https://doi.org/10.1080/037454809495210>
- Yefremova, Z.A., Ebrahimi E. & Yegorenkova, E. (2007) The subfamilies Eulophinae, Entedoninae and Tetrastichinae in Iran, with description of new species (Hymenoptera, Eulophidae). *Entomofauna*, 28 (30), 405–440.
- Yoder, M.J., Mikó, I., Seltmann, K.C., Bertone, M.A. & Deans, A.R. (2010) A gross anatomy ontology for Hymenoptera. *PLoS ONE*, 5 (12), 15991. <https://doi.org/10.1371/journal.pone.0015991>

زنبور *Diglyphus anadolucus* Doğanlar, 1982 (Hym., Eulophidae) پارازیتوئید خارجی و اولیه مرحله لاروی دوبالان خانواده Anthomyiidae، و فهرست گونه‌های جنس *Diglyphus* در ایران

یونس کریم‌پور<sup>\*</sup>، مجید جعفرلو<sup>۱</sup> و حسین لطفعلی‌زاده<sup>۲</sup>

۱ گروه گیاهپزشکی، دانشکده کشاورزی، دانشگاه ارومیه، ایران

۲ بخش تحقیقات رده‌بندی حشرات، موسسه تحقیقات گیاهپزشکی کشور، سازمان تحقیقات، آموزش و ترویج کشاورزی، تهران، ایران

\* پست الکترونیک نویسنده مسئول مکاتبه: [y.karimpour@urmia.ac.ir](mailto:y.karimpour@urmia.ac.ir)

ا تاریخ دریافت: ۰۳ دی ۱۴۰۱ | تاریخ پذیرش: ۰۹ بهمن ۱۴۰۱ | تاریخ انتشار: ۱۶ بهمن ۱۴۰۱

**چکیده:** این اولین مورد از ثبت میزبان برای زنبورهای پارازیتوئید جنس *Diglyphus* Walker (Hymenoptera, Eulophidae) از دوبالان خانواده Anthomyiidae است. این نتیجه، براساس نمونه‌هایی از زنبور *D. anadolucus* Doğanlar, 1982 بود که از پرورش لاروهای این زنبور روی لاروهای مگس *Pegomya terebrans* (Rondani) به عنوان مگس مینوز خارپنبه، (*Onopordum acanthium* L. (Asteraceae)) به دست آمدند. زنبور *D. anadolucus* برای اولین بار از ایران گزارش می‌شود. برخی نکات زیستی و ویژگی‌های ریخت‌شناسی این زنبور به همراه تصاویر مربوطه معرفی شد. فهرست به‌روز ۱۲ گونه از جنس *Diglyphus* و اطلاعات موجود از روابط میزبانی آنها در ایران نیز ارائه شد.

**واژگان کلیدی:** مهار زیستی، مینوز برگ، روابط میزبانی جدید، گزارش جدید